



## Loss of IL-10 secretion by regulatory B lymphocytes in multiple sclerosis patients

Laure Michel, Nicolas Degauque, Alexandra Garcia, Marion Salou, Annie Elong Ngono, Jean-Paul Soulillou, David Laplaud, Sophie Brouard

### ► To cite this version:

Laure Michel, Nicolas Degauque, Alexandra Garcia, Marion Salou, Annie Elong Ngono, et al.. Loss of IL-10 secretion by regulatory B lymphocytes in multiple sclerosis patients. 6th european workshop on immune-mediated inflammatory diseases, Nice, France. pp.P25. inserm-00643972

**HAL Id: inserm-00643972**

**<https://www.hal.inserm.fr/inserm-00643972>**

Submitted on 23 Nov 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



POSTER PRESENTATION

Open Access

# Loss of IL-10 secretion by regulatory B lymphocytes in multiple sclerosis patients

Laure Michel<sup>1,2\*</sup>, Nicolas Degauque<sup>1</sup>, Alexandra Garcia<sup>1</sup>, Marion Salou<sup>1</sup>, Annie Elong Ngono<sup>1</sup>, Jean-Paul Soullillou<sup>1</sup>, David Laplaud<sup>1,2</sup>, Sophie Brouard<sup>1</sup>

From 6th European Workshop on Immune-Mediated Inflammatory Diseases  
Nice, France. 23-25 November 2011

## Background

Recently, an alternative role of B cells has emerged. Regulatory B cells have been shown to down-regulate immune responses in mice and humans. These cells control disease progression in several models of human autoimmune diseases, including a model of Multiple Sclerosis (MS). Their regulatory function mostly relies on their capacity to produce IL-10.

## Objectives

Our aim was to explore the frequency, the phenotype and the functional properties of regulatory B cells in MS patients compared to Healthy Volunteers (HV).

## Methods

All patients suffered from MS and had not received immune drugs since at least 6 months. The frequency and the phenotype of B cell subsets have been analysed using different specific markers: CD19<sup>+</sup>CD27<sup>+</sup> memory B cells, CD19<sup>+</sup>CD38<sup>dim</sup>CD24<sup>dim</sup> mature naïve B cells and CD19<sup>+</sup>CD24<sup>high</sup>CD38<sup>high</sup> transitional B cells. Their capacity to secrete IL-10 was analysed *in vitro* 48 hours after stimulation by CD40 ligand and CpG ODN.

## Results

Thirty-eight MS patients (mean age: 41.55±12.8 yrs) and 21 HV (mean age: 35.4±13.4 yrs, NS) have been included. The patients had different MS forms: Clinically Isolated Syndrome (n=5), Relapsing-Remitting (n=23), Secondary Progressive (n=4) and Primary Progressive (n=6).

No significant difference was found for the frequency of the different B cell subsets in the different subgroups

of patients. Particularly, MS patients harbor the same number (in frequency and absolute value) of CD19<sup>+</sup>CD24<sup>high</sup>CD38<sup>high</sup> transitional B cells. However, the frequency of IL-10 secreting B cells was significantly decreased in MS patients (0.9±0.5%, n=9) compared to HV (2±0.95%, n=10, p<0.05, Mann Whitney test).

## Discussion/perspectives

MS patients display the same frequency of CD19<sup>+</sup>CD24<sup>high</sup>CD38<sup>high</sup> transitional B cells, described as having regulatory properties. Nevertheless, B cells of MS patients present a significant decreased secretion of IL-10, supporting a defect in the B cells regulatory property. If confirmed, these results could have a considerable impact for the development of new therapeutic strategies.

## Author details

<sup>1</sup>INSERM U643, Nantes University, Nantes, France. <sup>2</sup>Neurology Dept., Nantes Hospital, Nantes, France.

Published: 23 November 2011

doi:10.1186/1479-5876-9-S2-P25

Cite this article as: Michel et al.: Loss of IL-10 secretion by regulatory B lymphocytes in multiple sclerosis patients. *Journal of Translational Medicine* 2011 **9**(Suppl 2):P25.

<sup>1</sup>INSERM U643, Nantes University, Nantes, France  
Full list of author information is available at the end of the article